

Primary References Annotated Citations

- Austin MA, Breslow JL, Hennekens CH, Buring JE, Willett WC, Krauss RM. Low density lipoprotein subclass patterns and risk of myocardial infarction. *JAMA*. 1988;260:1917-1921.
In this case-controlled study, a preponderance of small, dense LDL particles correlated with a history of myocardial infarction, and was associated with the presence of both decreased HDL cholesterol and elevated triglyceride.
- Dreon DM, Fernstrom H, Miller B, Krauss RM. Low density lipoprotein subclass patterns and lipoprotein response to a reduced-fat diet in men. *FASEB J*. 994;8:121-126.
In subjects who reduced the caloric contribution of fat in their diets from a mean 41% to 25%, LDL-C and apo B were reduced significantly more in those with the small LDL trait at baseline than those without the trait. However, 41% of patients without the trait during the higher-fat diet manifested it on the lower-fat diet.
- Gardner CD, Fortmann SP, Krauss RM. Small low density lipoprotein particles are associated with the incidence of coronary artery disease in men and women. *JAMA* 1996;276:875-881.
LDL particle size correlated with CAD independently of and more strongly than levels of HDL-C, triglyceride, non-HDL cholesterol, triglyceride, smoking, and body mass index, although the ratio of the total cholesterol to HDL-C was a stronger independent predictor.
- Hecht HS, Superko HR. Electron beam tomography and National Cholesterol Education Program guidelines in asymptomatic women. *J Am Coll Cardiol*. 2001;37:1506-1511.
Among women with subclinical coronary atherosclerosis by electron beam computed tomography, 46.5% had LDL-C and HDL-C levels that were defined as low-risk for CAD by the guidelines of the 2nd adult treatment panel of the National Cholesterol Education Program (NCEP-II).

- Johansson J, Carlson LA, Landou C, Hamsten A. High density lipoproteins and coronary atherosclerosis. A strong inverse relation with largest particles is confined to normotriglyceridemic patients. *Am J Cardiol.*1991;11:174-182.
In men who underwent serial angiography after surviving a premature myocardial infarction, a higher distribution of HDL2b at the time of the follow-up angiogram correlated inversely with the severity and progression of CAD. This correlation was confined to a subset of patients with normal triglyceride levels.
- Krauss RM, Blanche PJ. Detection and quantitation of LDL subfractions. *Curr Opin Lipidol.* 1992;3:377-383, 1992.
This excellent review covers the use of ultracentrifugation and gradient gel electrophoresis for determining LDL particle size, identifying LDL subclasses, and measuring the distribution of these subclasses.
- Lamarche F, Tchernof A, Moorjani S, et al. Small, dense low-density lipoprotein particles as a predictor of the risk of ischemic heart disease in men. *Circulation.* 1997;95:69-75.
In a large prospective study, the small LDL trait predicted the future development of ischemic heart disease independently of the levels of LDL-C, HDL-C, apoprotein B, and, perhaps most importantly, triglycerides.
- Miller BD, Alderman EL, Haskell WL, Fair JM, Krauss RM. Predominance of dense low-density lipoprotein particles predicts angiographic benefit or therapy in the Stanford Coronary Risk Intervention project. *Circulation.*1996;94:2146-2153.
After four years of intensive treatment to improve risk factors in men with CAD, those with the small LDL trait at baseline had significantly greater angiographic regression of CAD than did those with predominantly large LDL particles.
- Skoglund-Andersson C, Tang R, Bond MG, de Faire U, et al. LDL particle size distribution is associated with carotid intima-media thickness in healthy 50-year-old men. *Arterioscler Thromb Vasc Biol* 1999;19:2422-2430.
The distribution of LDL IIIa + IIIb within the LDL spectrum correlates with the intima-media thickness of the carotid artery on B-mode ultrasound.
- Superko HR. Did Grandma give you heart disease? The new battle against coronary artery disease. *Am J Cardiol.* 1998;82(suppl):34Q-46Q.
A review of emerging risk factors in coronary heart disease, including the small LDL trait, elevated lipoprotein(a), and homocysteinemia.
- Superko HR. Lipid altering drugs LDL/HDL subclass distribution: Match the treatment to the disorder. *ACC Current Journal Review.* March/April, 2000:6-12.
This review discusses the identification and management of atherogenic patterns of LDL and HDL subclasses.

- Superko HR, Krauss RM. Differential effects of nicotinic acid in subjects with different LDL subclass patterns. *Atherosclerosis*. 1992;95:69-76.
Among patients at a high risk for CAD events, nicotinic acid produced a significantly greater decrease in LDL size in those with small LDL trait at baseline.
- Superko HR, Krauss RM, DiTicco C. Effect of HMGCoA reductase inhibitor (fluvastatin) on LDL peak particle diameter. *American J Cardiology*. 1997;80:78-81.
Fluvastatin has no effect on LDL size.
- Watts GF, Madalia S, Brunt JNH, Slavin BM, Coltart DJ, Lewis B. Independent associations between plasma lipoprotein subfraction levels and the course of coronary artery disease in the St. Thomas' Atherosclerosis Regression Study (STARS). *Metabolism*. 1993;42:1461-1467.
In men with CAD and hypercholesterolemia, reduction in LDL size was the most potent predictor of angiographic improvement among several lipoprotein variables.
- Williams PT, Krauss RM, Vranizan KM, Wood PD. Changes in lipoprotein subfractions during diet-induced and exercise-induced weight loss in moderately overweight men. *Circulation*. 1990;81:1293-1304.
Exercise- and diet-induced weight loss in overweight men are separately and significantly associated with improvement in HDL subclass distribution and an increase in LDL size.
- Williams PT, Superko HR, Alderman A. Small low density lipoprotein III, but not low density lipoprotein-cholesterol is related to arteriographic progression. *Circulation*. 2000;102(suppl):II-848 – II-849. Abstract 4076A
Arteriographic improvement during lipid-modifying therapy correlates with increases in the diameter of the most prevalent LDL particle (peak particle diameter) and with decreases of LDL distribution in regions IIIb and IVb.
- Zhao X, Kosinski AS, Malinow MR, Superko HR, King SB. Association of LDL heterogeneity and native coronary disease progression at 3 years following PTCA or CABG in EAST. *Circulation*. 2000;102(suppl):II-698 – II-699. Abstract 3378.
Arteriographic progression of CAD correlates with a high overall baseline distribution of LDL particles in the small regions (IIIa-IVb) and a small peak particle diameter.